



STATUS OF NOMADIC TRIBAL WOMEN IN NORTH KARNATAKA: IN TERMS OF ANTHROPOMETRY, HEALTH STATUS AND QUALITY OF LIFE

***Deeksha Naik, Kasturiba B. and Usha Malagi**

Department of Food Science and Nutrition, College of Rural Home Science,
University of Agricultural Sciences, Karnataka, India

ARTICLE INFO

Article History:

Received 14th February, 2018
Received in revised form
24th March, 2018
Accepted 07th April, 2018
Published online 31st May, 2018

Key Words:

Nomads, Tribe, Adult Women,
Anthropometry, Nomads,
Women, Nutritional status,
Quality of life.

ABSTRACT

Nomads are known as a group of communities who travel from place to place for their livelihood. The paper investigates Nutritional status of Nomadic Tribal women residing in North Karnataka. 30 nomadic adult women aged 18 to 45 years from 6 districts of North Karnataka were selected. Nutritional status in terms of Anthropometry was assessed. Results revealed that 40% of the women belong to underweight category of BMI. Poor nutritional status of women makes her more susceptible to infectious diseases. Ultimately it affects the productivity of women and in turn development of the community and the Nation.

Copyright © 2018, Deeksha Naik, Kasturiba B. and Usha Malagi. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Citation: Deeksha Naik, Kasturiba B. and Usha Malagi, 2018. "Status of nomadic tribal women in north karnataka: in terms of anthropometry, health status and quality of life", *International Journal of Development Research*, 8, (05), 20492-20494.

INTRODUCTION

The tribal population in India, though a numerically small minority, represents an enormous diversity of groups. They vary among themselves in respect of language and linguistic traits, ecological settings in which they live, physical features, size of the population, the extent of acculturation, dominant modes of making a livelihood, level of development and social stratification. They are also spread over the length and breadth of the country though their geographical distribution is far from uniform. Nomads are known as a group of communities who travel from place to place for their livelihood. Some are salt traders, fortune tellers, conjurers, ayurvedic healers, jugglers, acrobats, actors, story tellers, snake charmers, animal doctors, tattooists, grindstone makers, or basket makers. All told, anthropologists have identified about 5 nomadic groups in India, numbering perhaps 1 million people—around 1.2 percent of the country's billion-plus population. Women constitute almost 50% of world's total population.

In India this percentage is about 48.27, according to 2011 census. Country having such a high ratio of women section still lag behind in their status and empowerment. The situation is more severe among the tribal or primitive societies, which constitute approximately 8.9% of the total population of the country. India is also the second largest country to have highest concentration of indigenous population after the African continent. Lack of awareness about nutritional requirements mostly leaves the tribal women weak, anemic and they suffer from various diseases. Therefore in this study an effort is made to assess the nutritional status of the nomadic tribal adult women in North Karnataka.

MATERIALS AND METHODS

Nomadic tribal families migrating in North Karnataka were selected for the study. Six districts namely Dharwad, Hubli, Bijapur, Bagalkot, Gulbarga and Yadgiri of North Karnataka is the study location. A total of 30 nomadic tribal women of 18-45 years of age were selected for the study. Nutritional status was assessed by anthropometric measurements. The anthropometric measurements namely body weight and standing height were measured and recorded as per the

***Corresponding author: Deeksha Naik,**
Department of Food Science and Nutrition, College of Rural Home Science, University of Agricultural Sciences, Karnataka, India

guidelines suggested by Jelliff (1966). Weight was measured on the portable weighing square spring balance with light clothing and shoe removed and recorded to the nearest 0.5 kg. Standing height was measured with an anthropometric rod to the nearest of 0.1 cm. The subjects were made to remove the shoes and stand on the flat floor by the scale with feet parallel and heel, shoulder and back in upright posture, readings were recorded. Waist circumference and hip circumference of the individuals were also measured. BMI and WHR were computed for the data. Health problems such as head ache, leg pain, back pain, giddiness and leg cramp faced by these women were noted. Quality of life was assessed using WHOQoL-BREF scale. WHOQoL-BREF is an abbreviated generic Quality of Life Scale developed through the World Health Organization. Quality of life is defined as individuals perceptions of their position in life in the context of the culture and value systems in which they live and in relation to their goals, expectation, standards and concerns. The WHOQoL-BREF contains total of 26 questions. To provide a broad and comprehensive assessment, one item from each of the 24 facets contained in the WHOQoL-100 has been included. In addition, two items from the overall quality of life and general health facet have been included.

RESULTS AND DISCUSSION

The mean anthropometric measurements of adult women is represented in Table 1. Mean height was 153.86±7.3, mean weight was 47.23±5.6, mean waist circumference was 58.6±8.7, mean hip circumference was 76±11.5, mean WHR was 0.77±0.06 and mean BMI was 20.86±3.4 (Table-1) Classification of adult nomadic tribal women based on BMI is presented in Table 2. More than half of the subjects had ideal BMI (53.34%), nearly half of the subjects belonged to underweight category (40%), and about 3.33% subject belonged to overweight and obese grade 1 category. This reflects the availability of quality and quantity energy and other macro and micro nutrients and their physical activity (Devi and Sindhuja, 2015). Similar results were seen by Maiti (2005) that is 41% of both tribal and non tribal women had BMI of less than 18.5 kg/m². Classification of nomadic tribal women based on WHR is denoted in Table 3. Majority subjects belonged to no risk category (96.66%), only about 3.33% subject belonged to risky category. List of health problems faced by nomadic tribal women is presented in Table 4. Half of the subjects suffering from headache (53.33 per cent), followed by back pain (40 per cent), leg pain (26.66 per cent), giddiness (16.67 per cent) and leg cramp (13.33 per cent). Health problems such as head ache, back pain, leg pain, giddiness and leg cramp, might be because of their occupational activities and weakness. It was a striking feature that though the nomadic tribal women were suffering from high level of morbidity and mortality, they were not interested in going to hospitals. Majority of women were anemic. The reasons for undernutrition among tribal women could be poor dietary intake, ignorance, high morbidity due to unhygienic practices and surroundings. Similar results were observed in a study conducted by Maiti *et.al.* (2005). Classification of Quality of Life of nomadic tribal women according to domains is presented in Table 5. Entire population belonged to medium category under physical domain. Under physiological domain 13.33 per cent belong to high, 80.00 per cent to medium and 6.67 per cent to low category. Under social domain 26.67 per cent belong to high, 46.67 per cent to medium and 26.67 per

cent to low category. 20 per cent of population had high and 80 per cent had medium overall quality of life.

Table 1. Mean anthropometric measurements of Women N=30

Measurements	Mean
Weight (kg)	47.23 ± 5.6
Height (cm)	153.86 ± 7.3
Waist circumference (cm)	58.6 ± 8.70
Hip circumference (cm)	76 ± 11.50
Indices	
BMI	20.86 ± 3.40
WHR	0.77 ± 0.06

Table 2. Classification of adult nomadic tribal women based on BMI N=30

BMI classification	Presumptive diagnosis	No. of Nomadic women
< 18	Underweight	12 (40.00)
18.5-22.9	Ideal BMI	16 (53.34)
> 23	Over weight	1 (3.33)
> 25	Obese Grade I	1 (3.33)
> 30	Obese Grade II	-
Total		30 (100)
	Mean ± SD	20.86 ± 3.4

Figures in the parenthesis indicate percentages

Table 3. Classification of nomadic tribal women based on WHR N=30

Classification	No. of Nomadic women	%
No risk (<0.8)	29	96.66
At risk (>0.85)	1	3.33

Figures in the parenthesis indicate percentage

Table 4. List of health problems faced by nomadic tribal women

Health problems	Frequency	Percentage
Leg pain	8	26.66
Back pain	12	40.00
Head ache	16	53.33
Giddiness	5	16.67
Leg cramp	4	13.33



Women preparing food for her family

Table 5. Classification of Quality of Life of nomadic tribal women according to domains

	Physical domain		Physiological domain		Social relationships		Environment		Overall QOL	
	n	%	n	%	N	%	n	%	n	%
High	0	0.00	4	13.33	8	26.67	0	0.00	6.00	20.00
Medium	30	100	24	80.00	14	46.67	24	80.00	24.00	80.00
Low	0	0.00	2	6.67	8	26.67	6	20.00	0	0.00



Women participating in occupational activities along with men



Cleaning of utensils using dirty water



Roadside outdoor cooking of food

Conclusion

Poor nutritional status of women makes her more susceptible to infectious diseases. Under nutrition of mothers may be carried over to the children. Hence there is need to provide special attention to this group in improving their nutritional status by intervening appropriate health and nutritional programme like nutrition education, iron supplementation and deworming during all the different physiological stages of life. Ultimately it affects the productivity of women and in turn development of the community and the Nation, since in some states every fourth person is a tribe.

REFERENCE

Banik Datta Sudip., Bose Kaushik., Bisai Samiran., Bhattacharya Muthu., Das Subal., Jana Arpita and Purkait, 2007, Undernutrition among adult Dhimals of Naxalbari,

West Bengal: Comparison with other tribes of Eastern India, *Food and Nutrition bulletin*,.28(3):348-352.
 Bhasin.M.K and Jain Sweta, 2007, Biology of the tribal groups of Rajasthan, India: 1. Body mass index as an indicator of Nutritional status, *Anthropologist*, 9(3):165-175.
 Devi and Sindhuja, 2015, nutritional status and knowledge, life style ad dietary practices of tribal adult women, *Int.J of Recent Scientific Research*, 6(6),4449-4452.
 Maiti Sutapa., Unisa Sayeed and Agarwal K. Praween, 2005, Health care and health among Tribal women in Jharkhand: A situational Analysis, *Stud. Tribes Tribals*, 3(1):37-46.