



Full Length Research Article

**EFFECT OF MERCURIC CHLORIDE ON GILL STRUCTURE OF A FRESHWATER FISH
OREOCHROMIS MOSSAMBICUS (BLOCH)**

***Dr. Mukesh Kumar Napit**

Department of Zoology, Swami Vivkanand Government College Berasia, Bhopal (M.P.)

ARTICLE INFO

Article History:

Received 27th July, 2014
Received in revised form
25th August, 2014
Accepted 03rd September, 2014
Published online 25th October, 2014

Key words:

Mercuric chloride,
Histology, Gills,
O. mossambicus

Copyright © 2014 Dr. Mukesh Kumar Napit. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Fresh water fishes are economically important and an alternative source of significant human dietary constituents. The experimental fishes treated with lethal concentrations (1.00, 1.50, 0.63 and 2.00 ppm) of mercuric chloride showed many histological changes during 1 to 4 days of exposure. The gills showed vacuolization in the gill stem, gill lamellae ruptured, connective tissue cells in the stem damaged, destructed and congestion of haemocytes in the gill lamellae are observed. The present work on histological observation was carried out to know lesions in gills that had resulted from lethal exposure of the freshwater fish *O. mossambicus* to mercuric chloride.

INTRODUCTION

Histological studies have a way for understanding the pathological conditions of the animal by helping in diagnosing the abnormalities or damages of the tissues exposed to toxic stress of heavy metals (Sprague, 1971; Andhale, *et al.*, 2011, Maryam, *et al.*, 2013). Histological changes not only give an early indication of water pollution hazard, but also provide useful data on nature and degree of damage to cells and tissues (Shaikh, 2010). The gills of aquatic organism represent primary target of disturbance by pollutants as they are in direct contact with the external medium to perform gaseous exchanges and ionic regulations. This organ has linking sites to promote its regular functions, and these sites link to toxic substances with differentiated charges, triggering, mechanical responses and toxic effects to the organism (Jadhav, *et al.*, 2007). Allen (1995) observed the histological changes in gills of the *O. mossambicus* due to cadmium toxicity and observed ruptured gill lamellae. Histological changes in tissues are biomarkers of effect and exposure that integrate responses to contaminants at the cellular level. Vijayalaxmi and Tilak (1996) while working on *Labeo rohita* found pathological lesions in gill exposed to pesticides.

Saksena and Pandey (1993) also have observed toxic effect of copper on the histological effect on gills of a freshwater fish, *Labeo rohita*. In 1990, Victor *et al.*, observed the changes in the gill tissue of freshwater prawns, *Macrobrachium idea* exposed to mercury. A review of literature shows that there are more study is require to observed the histological changes caused by heavy metals in gills of *O. mossambicus*. Hence, the present study was undertaken to study the effects of mercuric chloride on histological changes in the gills of *O. mossambicus*

MATERIALS AND METHODS

Collection of fish The freshwater fishes, *O. mossambicus* were collected from the outskirts of Fresh water bodies of Bhopal city. They were acclimatized to laboratory conditions under normal day/night (11 L: 13 D) illumination at $27 \pm 1^\circ$ C for one week in plastic troughs (15" diameter) containing sufficient fresh water so that fishes are submerged. Before experimentation intermoult female fishes of approximately equal scales width (45-50 mm) and body weight (50 to 55 g) were sorted. For histological study fishes were split into 2 groups (Control, HgCl₂ treated), 5 fishes in each group and maintained under laboratory condition. After exposure period surviving fishes were sacrificed and the gills were quickly excised and utilized for histological studies from both the

***Corresponding author: Dr. Mukesh Kumar Napit**
Department of Zoology, Swami Vivkanand Govt. College Berasia,
Bhopal (M.P.)

control and experimental fishes. **Histological studies:** The excised tissues were fixed in aqueous Bouin's fluid. After fixation for 24 h the tissues were further processed to study histological details as per procedure of Munshi (1990). In brief the tissues were dehydrated through 30 - 100 % different alcohol grades and cleared in xylene. Cold and hot impregnations were followed by embedding the tissue in paraffin wax (M.P. 58-60 ° C). Serial sections were cut at 7 µm serial using rotary microtome. The section of gills was stained using Harris Haematoxylin and Eosin-Y as counter stain Munshi (1990). Damage to the tissues of treated fishes is recorded by comparing the data obtained from control.

RESULTS

The histological study has been carried out to know lesions in gills that had resulted from lethal exposure of the freshwater female fishes, *O. mossambicus* to mercuric chloride. In freshwater fish gills are paired and have crescent shaped structure and located laterally on either side of the hepatopancreas and alimentary canal within the gill chamber in the antero-ventral region of thoracic region. Gills are normally deep brown in colour.

Histological structure of the gills in control fish

The gills of *O. mossambicus* are phyllobranchiate type consisting of central stem (axis, raphe) that bears serially the paired plate or leaf like lamellae, which are actually flattened sacs. The central axis has afferent and efferent haemal channels on each end. Three types of cells are found: (1) *Connective tissue cells* are located in the gill stem; (2) *Branched arthrocytes* are located in the stem and proximal lamellae; (3) *Lamellar cells* are located in the epithelium. Each gill plate has outer cuticular layer enclosing a single layer of gill epithelial cells. The gill epithelium stained deep pink in haematoxylin and eosin. The central axis contained large amount of darkly stained haemocytes. The central axis also had few nephrocytes. The nephrocytes are vacuolated large cells that have some amount of pale brown material (Fig. 1-A).




Figure 1a Crab Fish. Control crab x 400: Distinct haemocytes, gill stem, gill lamella within connective fluidy bands are observed

Histological changes in the gills of experimental fish

The experimental fishes treated with lethal concentration of mercuric chloride showed many histological changes during 1

to 4 days of exposure. In the experimental fish gills showed vacuolization in the gill stem, gill lamellae ruptured, connective tissue cells in the stem damaged, destructed and congestion of haemocytes in the gill lamellae are observed. Pillar cells are damaged. The thin connective fluidy band present in between the two gill lamellae is found ruptured (Fig. 1B).



Figure 1b Crab Fish. Experimental crab x 200: Vacuolization of gill stem, fusion and ruptured gill lamella, destructed and congestion of haemocytes, thin connective fluidy bands are ruptured

DISCUSSION

The experimental fishes exposed to lethal concentrations of mercuric chloride exhibit histological changes in the gills due to the accumulation of mercuric chloride within the organism body at lethal and levels lead to histological lesion in the body fish, *O. mossambicus*. Histopathological studies are also useful in evaluating the pollution potential of heavy metalpesticides, since trace amount of these chemicals which do not bring animal mortality over a given period, were capable of producing considerable organ damage (Kumar and Pant, 1984; Jaykumar, 2002). Despite much information available on the histological changes caused by heavy metalpesticides, the mode of action on the vital organs is still not fully understood. The difference between the control and the experimental tissues were studied critically. The study of micro-anatomy (histology) of the specific tissue constitutes an important diagnostic tool to observe the histological effects caused by a pollutant. The histological changes may be the manifestation of sick tissue (Kamble and Potdar, 2010). Gills of the fish, *O. mossambicus* were of phyllobranchiate type with a central axis and gill lamellae arranged in two rows on either side (Fig.-1 A and B). The structure of the gills observed in the present study was well in accordance with earlier reports on different fishes (Joshi, 2006; Jadhav, *et al.*, 2007).

In experimental fish the histological sections of gills showed vacuolization in the gill stem, gill lamellae ruptured, connective tissue cells in the stem damaged, destructed and congestion of haemocytes in the gill lamellae are observed. Pillar cells are damaged. The thin connective fluidy band present in between the two gill lamellae is found ruptured (Fig. 1-B). Li *et al.*, (2007) observed structural changes including the swelling and fusion of the lamellae; abnormal gill tips; and necrotic lamellae in gills of a giant freshwater prawn, *Macrobrachium rosenbergii* exposed to waterborne copper.

Kurian and Radhakrishnan (2002) observed such changes in the gills of the field fish, *Labeo rohita* on exposure to nickel. The histological techniques are the promising area of research in aquatic toxicology as it gives the real picture of the effects imposed and the involvement of the xenobiotics in either disturbing or destroying the vital organs of living organisms. Many researchers have reported the degenerative changes in selected tissues of the animals in response to pollution by various toxicants (Shanmugam *et al.*, 2000; Suresh, 2001; Kale, 2002; Reddy, 2005; Tilak, *et al.*, 2005; Andhale, *et al.*, 2011, Mukke, 2012).

Conclusion

The histology is a most common tool for determining the deleterious effects of toxic substances on the treated animals. In the present investigation an attempt has been made to evaluate the intensity of the damage done to gills of crab subjected to its lethal concentrations of mercuric chloride.

Acknowledgements

Author Napit M.K. is thankful to University Grant Commission (UGC), Regional Office Bhopal for financial assistance through Minor Research Project (MRP) during this research work. Also thankful to Office Staff of Govt. College Berasia Bhopal for valuable support.

REFERENCES

- Allen R. 1995, Observed the Histochemical changes in gill of the *O. mossambicus*. *Int. J. of Global Bioscience*. vol.2(3) 2013, pp212-217
- Bancroft, J.D. and Stevens, A. 1982: Theory and Practice of Histological Techniques. *Churchill Livingstone. Publ.* New York.
- Jadhav, T.J., Kulkarni, G.K., Joshi, P.P. and Jawale, C.J. 2007: Effect of endosulfan and thimet on histological profile of gills and ovary of a freshwater fishes *O. mossambicus* In Aquaculture: B.N. Pandey, Deshmukh, S. and P.N. Pandey Eds. *A.P.H. Publ. Corp.* New Delhi: 85-92.
- Diwan, A.D. 1973: Studies on sexual maturity and growth rate studies of the freshwater fish *O. mossambicus* Marath. *Univ. J. Sci.* 12:125.
- Joshi, P.P. 2006: Physiological responses of freshwater fish and crab to pyrethroid pesticides: A comparative study. Ph.D. Thesis submitted to Dr. Babasaheb Ambedkar Marathwada University, Aurangabad.
- Kale, M.K. 2002: Cadmium toxicity to some physiological activities of freshwater fish and crab: A comparative study. Ph.D. Thesis submitted to Dr. Babasaheb Ambedkar Marathwada University, Aurangabad.
- Sekar, P. 2005: Impact of textile dye industry effluent on a freshwater field female fishes *O. mossambicus* (Herbst.). A Ph.D. Thesis submitted to the University of Madras.
- Suresh, V. 2001: A study on the effects of heavy metals toxicity on a brackish water fishes, *Uca lacteal annulipes* (Crane, 1975) of Pulicat Lake, TN, Ph.D. Thesis. University of Madras, Chennai.
- Andhale, A.V., Bhosale, P.A. and Zambare, S.P. 2011: Histopathological study of nickel induced alterations in the fresh water bivalve, *Lammellidens marginalis*. *Journal of Experimental Sciences* Vol. 2, Issue 4: 01-03.
- Kamble, N.A. and Potdar, V.V. 2010: Hepatopancreatic damages in snail *Bellamya bengalensis* (L) tested against lead acetate toxicity. *Indian Journal of Comparative Animal Physiology*. Vol. 28(1): 7-10.
- Kumar, S and Pant, S.C. 1984: Organal damage caused by aldrin to a freshwater teleost *Barbus conchonus* Hamilton. *Bull. Environ. Contam. Toxicol.* 33: 50-55.
- Kurian, M.A. and Radhakrishnan T. 2002: Study on gill of field crab, *Paratelphusa hydrodromus* (Herbst.) exposed to nickel. *J. Environ. Biol.*, 23:151-155.
- Li, N., Zhao, Y. and Yang, J. 2007: Impact of waterborne copper on the structure of gills and hepatopancreas and its impact on the content of metallothionein in juvenile giant freshwater prawn, *Macrobrachium rosenbergii* (Crustacea: Decapoda). *Arch. Environ. Contam. Toxicol.* 52:73-79.
- Mazon, A.F., Nolan, D.T., Lock, R.A.C., Fernandes, M.N. and Wendelaar Bonga, S.E. 2004: A short-term in vitro gill culture system to study the effects of toxic (copper) and non-toxic (cortisol) stressors on the rainbow trout, *Oreochromis mossambicus* (Walbaum). *Toxicol in Vitro* 18:691-701.
- Mukke, V. K. 2012: Bioaccumulation study of mercury chloride in selected tissue of fresh water fishes *O. mossambicus* from Aurangabad region. *Journal of Chemical and Pharmaceutical Research*, 4(1):398-401.
- Richmonds, C. and Datta, H.M. 1989: Histopathological changes induced by malathion in the gills of *Lepomis macrochirus*. *J. Bull. Environ. Contam. Toxicol.* 43: 123-136.
- Saksena, D.N. and Pandey, R. 1993: Copper Sulphate induced histopathological alteration in the gill of fingerlings of an Indian major carp, *Labeo rohita* (Hamilton). *J. Natcon.* 5(1):65-69.
- Sarojini, R. Machale, P.R. and Nagabhushanam, R. 1990: Adverse effect of zinc sulphate exposure to ovarian development of freshwater crab, *Barytelphusa guerini*. *J. Natcon.* 2(2):97-104.
- Shanmugam, M., Venkateshwarlu, M. and Naveed, A. 2000: Effect of pesticides on the Freshwater fishes *O. mossambicus* (Westwood). *J. Ecotoxicol. Environ. Monit.* 10:(273-279).
- Shaikh, F.I, Ustad, I.R. and Ansari, N.T. 2010: Effect of heavy metal on the ovary of freshwater fishes *O. mossambicus* (Westwood). *The Bioscan*, 5(2): 335-338.
- Sprague, J. B. 1971: The measurement of pollutant toxicity to fish III, sub lethal effect and safe concentration. *Water Res.* 5:245-266.
- Tilak, K.S., Veeraiah, K. and Thathaji, P.B. 2005: Histopathological changes in gill tissues of the fish *Channa punctatus* exposed to sublethal concentration of butachlor and machete. *J. Aqua. Biol.* 20(1):111-115.
- Vernberg, W. B. and Vernberg, F. J. 1972: The synergistic effect of temperature salinity and mercury on survival and metabolism of the adult fiddler fishes. *Fish Bull.* 70: 415-420.
- Victor, B., Narayanan, M. and Nelson, D.J. 1990: Gill pathology and hemocytes response in mercury exposed *Macrobrachium idae* (Heller). *J Environ Biol* 11: 61-65.
- Vijayalakshmi, S. and Tilak, K.S. 1996: Effect of pesticides on the gill morphology of *Labeo rohita*. *J. Ecotoxicol. Environ. Monit.* 6(1): 59-64.
- Wu, J.P., Chen, H.C. and Huang, D.J. 2008: Histopathological and biochemical evidence of hepatopancreatic toxicity caused by cadmium and zinc in the white shrimp, *Litopenaeus vannamei*. *Chemosphere* 73: 1019-1026.
- Reddy, A.P. 2005: Histopathological changes observed in the gill, liver and kidney of fish *Labeo rohita* exposed to confidor. Proc. International Conf. on Environmental Management. Institute of Science and Technology, Centre for Environment, Hyderabad. 28-30, October.