ISSN: 2230-9926

Available online at http://www.journalijdr.com

IJDR

International Journal of Development Research
Vol. 16, Issue, 10, pp.71152-71154, October, 2026
https://doi.org/10.37118/ijdr.31254.10.2026



RESEARCH ARTICLE OPEN ACCESS

PREPARATION OF MOTHER TINCTURES IN HOMOEOPATHIC PHARMACY AND THEIR TRADITIONAL HOMOEOPATHIC INDICATIONS

Dr. Rizwan Ahmad (BHMS, MD), Miss. Kavita Adhagale and Mr. Faizan Ansari

Department of Homoeopathic Pharmacy, Institution, SPHMC & H Shrirampur

ARTICLE INFO

Article History:
Received 24th July, 2026
Received in revised form
21st August, 2026
Accepted 23rd September, 2026
Published online 31st October, 2026

Key Words:

Mother Tincture, Homoeopathic Pharmacy, Maceration, Percolation, Menstruum, Quality Control.

Corresponding author:
Dr. Rizwan Ahmad (BHMS, MD)

ABSTRACT

Mother tinctures are important stock preparations in homoeopathic pharmacy. They are generally prepared from fresh or dried plant materials, and in some cases other source materials, by extracting soluble constituents with a suitable menstruum such as alcohol, purified water, or a specified mixture. The exact method, drug strength, solvent composition, extraction time, and quality tests depend on the individual monograph in the applicable Homoeopathic Pharmacopoeia. This review describes the concept of mother tinctures, their principal preparation methods, essential quality-control measures, storage, and examples of traditional homoeopathic indications. The indications listed are descriptions of homoeopathic practice and materia medica—not proof that these products treat or cure the named diseases. Reliable clinical evidence supporting homeopathy for specific conditions remains limited, and mother tinctures may contain pharmacologically active substances and alcohol.

Copyright©2026, Rizwan Ahmed et al. *This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.*

Citation: Dr. Rizwan Ahmad (BHMS, MD), Miss. Kavita Adhagale and Mr. Faizan Ansari, 2026. "Preparation of mother tinctures in homoeopathic pharmacy and their traditional homoeopathic indications.". *International Journal of Development Research*, 16, (10), 71152-71154.

INTRODUCTION

A mother tincture (commonly designated by the symbol φ or Q) is an initial liquid preparation made from a source material according to a prescribed pharmacopoeial procedure. It may be used as a starting material for subsequent potentisation or, in certain preparations, dispensed as a tincture under professional supervision. Unlike highly diluted potencies, mother tinctures can contain measurable quantities of plant constituents. Consequently, accurate identification, controlled extraction, correct labelling, and safety assessment are essential.

OBJECTIVES

- To explain the meaning and pharmaceutical importance of mother tinctures.
- To outline the main methods used for their preparation.
- To describe basic quality-control, packaging, and storage requirements.
- To summarise examples of traditional homoeopathic indications while distinguishing traditional use from proven efficacy.

PRINCIPLE OF PREPARATION: Preparation is based on extraction: a suitable menstruum penetrates the processed raw material and dissolves extractable constituents. The resulting liquid is separated from the marc (spent plant material), clarified when required, and adjusted or processed as directed by the relevant monograph.

There is no single universal formula for all mother tinctures. Drug-to-menstruum ratio, alcohol percentage, plant moisture correction, and processing conditions must be taken from the official monograph rather than guessed or generalised.

MATERIALS AND EQUIPMENT

- Authenticated and correctly identified raw material, with batch and source records.
- Menstruum of pharmacopoeial quality (for example, specified-strength ethanol and/or purified water).
- Clean, suitable glass or stainless-steel extraction vessel; lid; weighing and measuring equipment.
- Knife or cutter / grinder as specified, filter or press, and suitable receiving vessel.
- Clean, compatible amber-glass storage bottles with tight closures and appropriate labels.
- Personal protective equipment and documented cleaning, hygiene, and safety procedures.

METHODS OF PREPARATION

Maceration: Maceration involves keeping the suitably processed raw material in the prescribed menstruum for a specified period. The vessel is closed and kept under the conditions stated in the

Calendula officinalis Q	Traditionally used externally in homoeopathic practice for minor superficial cuts and skin care.	External products only as labelled; avoid deep wounds, serious burns, and eye contact.
Hamamelis virginiana Q	Traditionally associated with bruised soreness and venous congestion in homoeopathic materia medica.	Persistent bleeding, painful swelling, or suspected thrombosis needs medical assessment.
Berberis vulgaris Q	Traditionally associated with urinary complaints and certain radiating pains in materia medica.	Flank pain, fever, blood in urine, or suspected kidney stones requires prompt medical evaluation.
Chelidonium majus Q	Traditionally associated with digestive and right-upper-abdominal complaints in materia medica.	Liver injury has been reported with Chelidonium-containing products; avoid unsupervised use.
Crataegus oxyacantha Q	Traditionally associated with cardiac symptoms in homoeopathic practice.	Not a substitute for evidence-based cardiac care; chest pain or breathlessness may be an emergency.
Passiflora incarnata Q	Traditionally associated with restlessness and sleep-related complaints in homoeopathic practice.	May interact with sedatives; persistent insomnia or severe anxiety warrants clinical assessment.
Echinacea angustifolia Q	Traditionally used in homoeopathic practice in relation to recurrent infections and inflammatory complaints.	Allergy is possible, especially in people sensitive to plants in the Asteraceae family.
Sabal serrulata Q	Traditionally associated with urinary symptoms attributed to prostate enlargement in homoeopathic materia medica.	Urinary retention, fever, blood in urine, or new urinary symptoms need medical assessment.
Urtica urens Q	Traditionally associated with itching or nettle-like skin eruptions in materia medica.	A rapidly spreading rash, facial swelling, or breathing difficulty requires urgent care.
Avena sativa Q	Traditionally associated with nervous exhaustion and convalescence in homoeopathic practice.	Persistent fatigue may have an underlying medical cause and should be assessed.

monograph; mixing or agitation is performed when required. The liquid is then separated from the marc, commonly by filtration or pressing, and further processed according to the official procedure. Maceration is useful where the material requires prolonged contact with the solvent or is not well suited to percolation.

Percolation: In percolation, prepared material is packed into a suitable percolator and the menstruum passes slowly through it. The rate of flow, wetting or imbibition period, solvent quantity, and collection procedure are controlled according to the monograph. The percolate is collected and processed as specified. Correct packing is important: overly tight packing can impede flow, while loose packing may allow channeling and incomplete extraction.

Other procedures: Some monographs may prescribe special procedures based on the nature of the starting material, including expression of fresh plant juice, adjustment for natural moisture, or other defined extraction steps. The official monograph takes precedence over a generic description.

GENERAL WORKFLOW

- Authenticate the source material and inspect it for identity, purity, and contamination.
- Clean and process the material as specified; determine moisture content where required.
- Calculate the exact amount and composition of menstruum from the relevant monograph.
- Perform maceration, percolation, or the specified special procedure under controlled conditions.
- Separate the liquid extract from the marc and complete any prescribed clarification or adjustment.
- Carry out applicable in-process and finished-product checks; document results and batch details.
- Fill into suitable clean containers, close securely, label correctly, and store as directed.

QUALITY CONTROL AND STORAGE

Quality assurance begins with authenticated raw materials and continues through extraction, filling, and storage. Depending on the monograph and applicable regulations, checks may include organoleptic characteristics, identity, specific gravity, alcohol content, dry residue or extractive value, and relevant chemical or microbiological tests. Manufacturing records should document raw-material lots, calculations, equipment, process conditions, yields, test results, deviations, and release decisions. Mother tinctures are generally stored in well-closed, compatible containers, protected from direct light and excessive heat, under the conditions specified by the applicable standard.

EXAMPLES OF TRADITIONAL HOMOEOPATHIC INDICATIONS

The following are illustrative descriptions commonly found in homoeopathic materia medica and practice. They are not validated treatment recommendations, and they should not be interpreted as evidence that a tincture is effective for a disease. Selection, formulation, route, and dose depend on the product monograph and qualified clinical judgement.

SAFETY, ETHICS, AND CLINICAL LIMITATIONS

Mother tinctures are not necessarily chemically inert: they may contain active plant constituents and significant amounts of alcohol, and some ingredients can cause adverse effects or interact with conventional medicines. Product quality may vary if raw materials, extraction, or manufacturing controls are inadequate. Preparation should be carried out only in a properly equipped, authorised pharmacy or laboratory by trained personnel, using the applicable official pharmacopoeial monograph and local regulations.

This article does not provide a do-it-yourself formula or dosing advice. Homeopathy has limited reliable evidence of effectiveness for treating specific diseases. Mother tinctures must not replace proven treatment, delay diagnosis, or be used as a substitute for emergency care. Seek qualified medical care for severe, persistent, or worsening symptoms. Extra caution is required for children, pregnancy, liver or kidney disease, alcohol sensitivity, and people taking other medicines.

CONCLUSION

Mother tinctures are foundational preparations in homoeopathic pharmacy and require accurate raw-material identification, monograph-based extraction, validated quality checks, appropriate packaging, and controlled storage. Maceration and percolation are major preparation methods, but their detailed parameters differ by source material and official standard.

Traditional indications should be presented accurately as descriptions of homoeopathic practice rather than as established clinical efficacy. Pharmaceutical quality, patient safety, and timely evidence-based medical care remain essential.

REFERENCES

Central Council for Research in Homoeopathy (CCRH). Indian Journal of Research in Homoeopathy. “Indian standards for methods of preparation of homoeopathic mother tinctures.” 2026;20(2):294–300. https://www.ijrh.org/ journal/ vol20/iss 2/18/ of AYUSH / Pharmacopoeia Commission for Indian Medicine & Homoeopathy. Homoeopathic Pharmacopoeia o

f India. Available at: https://pcimh.gov.in/WriteReadData/RTF1984/HPISeventhedition.pdf

National Center for Complementary and Integrative Health (NCCIH). Homeopathy: What You Need to Know. https://www.nccih.nih.gov/health/homeopathy

****